



CLPB deficiency

CLPB deficiency is a rare disorder characterized by neurological problems and a shortage of infection-fighting white blood cells (neutropenia). Signs and symptoms of the condition develop by early childhood, and their severity varies widely among affected individuals.

In the most severely affected individuals, features of CLPB deficiency are apparent in infancy and sometimes at birth. Affected babies have serious neurological problems, which can include an exaggerated startle reaction (hyperekplexia) to unexpected stimuli such as loud noises, reduced movement, muscle tone that is either decreased (hypotonia) or increased (hypertonia), swallowing problems, difficulty breathing, and recurrent seizures (epilepsy). These babies may also have movement abnormalities, such as difficulty coordinating movements (ataxia), involuntary tensing of the muscles (dystonia), or uncontrolled movements of the body (dyskinesia). In addition, these babies have recurrent, life-threatening infections due to severe neutropenia. Affected individuals are at risk of developing a blood cell disorder called myelodysplastic syndrome or a form of blood cancer called leukemia. Because of their severe health problems, affected infants usually live only a few weeks or months.

Moderately affected individuals have neurological problems similar to those described above, although they are less severe. They include hypotonia, muscle stiffness (spasticity), and movement abnormalities. Other features of moderate CLPB deficiency include epilepsy and mild to severe intellectual disability. Neutropenia in these individuals can lead to recurrent infections, although they are not life-threatening.

Mildly affected individuals have no neurological problems, and although they have neutropenia, it does not increase the risk of infections. Some people with mild CLPB deficiency develop deposits of calcium in the kidneys (nephrocalcinosis) or kidney (renal) cysts.

Many people with mild, moderate, or severe CLPB deficiency have clouding of the lenses of the eyes (cataracts) from birth (congenital) or beginning in infancy.

CLPB deficiency is associated with increased levels of a substance called 3-methylglutaconic acid in the urine (3-methylglutaconic aciduria). This abnormality, which provides a clue to the diagnosis, does not appear to cause any health problems.

Frequency

CLPB deficiency is a rare disorder; the prevalence is not known. At least 26 cases have been reported in the medical literature.

Genetic Changes

CLPB deficiency is caused by mutations in the *CLPB* gene, which provides instructions for making a protein whose function is unknown. Based on its similarity to a protein in other organisms, the CLPB protein is thought to help unfold misfolded proteins so they can be refolded correctly. If not fixed, misfolded proteins cannot function properly and may be damaging to cells.

CLPB gene mutations likely reduce or eliminate the amount of functional CLPB protein. The severity of the condition is thought to be related to the amount of functional protein remaining: severe CLPB deficiency is likely caused by a complete absence of CLPB protein, while moderate and mild CLPB deficiency result when some functional CLPB protein is produced. Researchers are unsure how reduction or absence of this protein leads to the signs and symptoms of CLPB deficiency.

Inheritance Pattern

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- 3-methylglutaconic aciduria-cataract-neurologic involvement-neutropenia syndrome
- 3-methylglutaconic aciduria type 7
- 3-methylglutaconic aciduria type VII
- 3-methylglutaconic aciduria with cataracts, neurologic involvement and neutropenia
- MEGCANN
- MGA7
- MGCA7

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: 3-methylglutaconic aciduria with cataracts, neurologic involvement, and neutropenia
<https://www.ncbi.nlm.nih.gov/gtr/conditions/CN228597/>

Other Diagnosis and Management Resources

- GeneReview: CLPB Deficiency
<https://www.ncbi.nlm.nih.gov/books/NBK396257>
- Seattle Children's Hospital: Neutropenia Symptoms and Diagnosis
<http://www.seattlechildrens.org/medical-conditions/heart-blood-conditions/neutropenia-symptoms/>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: Neutropenia - Infants
<https://medlineplus.gov/ency/article/007230.htm>
- Health Topic: Blood Disorders
<https://medlineplus.gov/blooddisorders.html>
- Health Topic: Movement Disorders
<https://medlineplus.gov/movementdisorders.html>

Additional NIH Resources

- National Eye Institute: Facts About Cataract
https://nei.nih.gov/health/cataract/cataract_facts

Educational Resources

- MalaCards: 3-methylglutaconic aciduria, type vii, with cataracts, neurologic involvement and neutropenia
http://www.malacards.org/card/3_methylglutaconic_aciduria_type_vii_wit_h_cataracts_neurologic_involvement_and_neutropenia
- Merck Manual Consumer Version: Neutropenia
<http://www.merckmanuals.com/home/blood-disorders/white-blood-cell-disorders/neutropenia>
- Orphanet: 3-methylglutaconic aciduria type 7
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=445038
- The University of Arizona Health Sciences: 3-methylglutaconic Aciduria with Cataracts, Neurologic Involvement, and Neutropenia
<http://disorders.eyes.arizona.edu/category/alternate-names/3-methylglutaconic-aciduria-type-vii>

Patient Support and Advocacy Resources

- Child Neurology Foundation
<http://www.childneurologyfoundation.org>
- CLIMB: Children Living with Inherited Metabolic Disorders
<http://www.climb.org.uk/>
- Immune Deficiency Foundation
<http://primaryimmune.org/>

GeneReviews

- CLPB Deficiency
<https://www.ncbi.nlm.nih.gov/books/NBK396257>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28CLPB+deficiency%29+OR+%283-methylglutaconic+aciduria+with+cataracts,+neurologic+involvement,+and+neutropenia%29+OR+%28CLPB+mutation%29+OR+%28CLPB+variants%29+OR+%28CLPB+disruption%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- 3-METHYLGLUTACONIC ACIDURIA WITH CATARACTS, NEUROLOGIC INVOLVEMENT, AND NEUTROPENIA
<http://omim.org/entry/616271>

Sources for This Summary

- Capo-Chichi JM, Boissel S, Brustein E, Pickles S, Fallet-Bianco C, Nassif C, Patry L, Dobrzeniecka S, Liao M, Labuda D, Samuels ME, Hamdan FF, Vande Velde C, Rouleau GA, Drapeau P, Michaud JL. Disruption of CLPB is associated with congenital microcephaly, severe encephalopathy and 3-methylglutaconic aciduria. *J Med Genet*. 2015 May;52(5):303-11. doi: 10.1136/jmedgenet-2014-102952.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25650066>
- GeneReview: CLPB Deficiency
<https://www.ncbi.nlm.nih.gov/books/NBK396257>
- Kanabus M, Shahni R, Saldanha JW, Murphy E, Plagnol V, Hoff WV, Heales S, Rahman S. Bi-allelic CLPB mutations cause cataract, renal cysts, nephrocalcinosis and 3-methylglutaconic aciduria, a novel disorder of mitochondrial protein disaggregation. *J Inher Metab Dis*. 2015 Mar; 38(2):211-9. doi: 10.1007/s10545-015-9813-0.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25595726>
- Saunders C, Smith L, Wibrand F, Ravn K, Bross P, Thiffault I, Christensen M, Atherton A, Farrow E, Miller N, Kingsmore SF, Ostergaard E. CLPB variants associated with autosomal-recessive mitochondrial disorder with cataract, neutropenia, epilepsy, and methylglutaconic aciduria. *Am J Hum Genet*. 2015 Feb 5;96(2):258-65. doi: 10.1016/j.ajhg.2014.12.020.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25597511>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4320254/>
- Wortmann SB, Zietkiewicz S, Kousi M, Szklarczyk R, Haack TB, Gersting SW, Muntau AC, Rakovic A, Renkema GH, Rodenburg RJ, Strom TM, Meitinger T, Rubio-Gozalbo ME, Chrusciel E, Distelmaier F, Golzio C, Jansen JH, van Karnebeek C, Lillquist Y, Lücke T, Öunap K, Zordania R, Yapfite-Lee J, van Bokhoven H, Spelbrink JN, Vaz FM, Pras-Raves M, Ploski R, Pronicka E, Klein C, Willemsen MA, de Brouwer AP, Prokisch H, Katsanis N, Wevers RA. CLPB mutations cause 3-methylglutaconic aciduria, progressive brain atrophy, intellectual disability, congenital neutropenia, cataracts, movement disorder. *Am J Hum Genet*. 2015 Feb 5;96(2):245-57. doi: 10.1016/j.ajhg.2014.12.013.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25597510>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4320260/>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/clpb-deficiency>

Reviewed: February 2017

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services